

The Fourier method for the linearized Davey-Stewartson I equation

O. M. Kiselev

Ufa Institute of Mathematics, Russian Academy of Sciences,
112 Chernyshevsky str., Ufa, 450000, Russia
E-mail: ok@imat.rb.ru

Abstract

The linearized Davey-Stewartson equation with varying coefficients is solved by Fourier method. The approach uses the inverse scattering transform for the Davey-Stewartson equation.

1 Introduction

The Davey-Stewartson equation (DS) is the well-known subject for investigations because of two causes. First, the equation models a nonlinear interaction between a long surface wave and a short surface wave [1]. Second, DS is integrable by an inverse scattering transform (IST) [2] and then one can investigate a solution structure in details.

In this work we consider a linear system of equations with varying coefficients, which come out when one linearizes the DS at a nonzero solution as a background. We develop the Fourier method for the above linear system. This method is based on IST results for DS [2, 3, 4]. The basic functions, which are used for the Fourier expansion, are associated with a scattering problem for a Dirac system.

A realized approach one can see as generalizing on (2+1)-dimensions (two spatial variables and time) system of pioneer works of D.Kaup [5, 6], in which the Fourier method was formulated for linearized (1+1)-dimensional integrable equations. The obtained results make possible to study perturbations of the DS, in general putting out the equation from the class of integrable equations.

2 The Dirac equation and the basic functions

Here we consider a Goursat problem for the Dirac equation [2]-[4]:

$$\begin{pmatrix} \partial_\xi & 0 \\ 0 & \partial_\eta \end{pmatrix} \psi = -\frac{1}{2} \begin{pmatrix} 0 & q_1 \\ q_2 & 0 \end{pmatrix} \psi \quad (1)$$

Define a solutions of (1) as follows [2]:

$$\begin{aligned} \psi_{11}^+|_{\xi \rightarrow -\infty} &= \exp(ik\eta), & \psi_{12}^+|_{\xi \rightarrow -\infty} &= 0, \\ \psi_{21}^+|_{\eta \rightarrow -\infty} &= 0, & \psi_{22}^+|_{\eta \rightarrow -\infty} &= \exp(-ik\xi); \\ \psi_{11}^-|_{\xi \rightarrow -\infty} &= \exp(ik\eta), & \psi_{12}^-|_{\xi \rightarrow +\infty} &= 0, \\ \psi_{21}^-|_{\eta \rightarrow -\infty} &= 0, & \psi_{22}^-|_{\eta \rightarrow -\infty} &= \exp(-ik\xi). \end{aligned}$$

Denote by $(\chi, \mu)_q$ a bilinear form:

$$(\chi, \mu)_q = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\xi d\eta (\chi_1 \mu_1 q_2 + \chi_2 \mu_2 q_1), \quad (2)$$

where χ_i and μ_i are elements of the columns χ and μ .

Denote by $\phi^{(1)}$ and $\phi^{(2)}$ the solutions which are dual to $\psi_{(1)}$ (first column of the matrix ψ^+) and $\psi_{(2)}$ (second column of the matrix ψ^-) with respect to the bilinear form (2).

It is known, that the solution of (1) satisfies a nonlocal Riemann-Hilbert problem [3]. Formulate this problem for ψ_{11}^- and ψ_{12}^+ . Denote by $\psi^{(1)}$ a row $\{\psi_{11}^-, \psi_{12}^+\}$, then [3]:

$$\psi^{(1)} = E^{(1)}(ik\eta) + S[s]\psi^{(1)}.$$

Here $E^{(1)}(z)$ is the first row of a matrix $E(z) = \text{diag}(\exp(z), \exp(-z))$, the operator $S[s]$ is defined by formula:

$$\begin{aligned} S[s]\psi^{(1)} &= \left[\exp(ik\eta) \left(\exp(-ik\eta) \int_{-\infty}^{\infty} dl s_1(k, l) \psi_{12}^+(\xi, \eta, l) \right)^- , \right. \\ &\quad \left. \exp(ik\xi) \left(\exp(-ik\xi) \int_{-\infty}^{\infty} dl s_1(k, l) \psi_{11}^-(\xi, \eta, l) \right)^+ \right], \end{aligned}$$

where

$$\left(f(k) \right)^{\pm} = \frac{1}{2i\pi} \int_{-\infty}^{\infty} \frac{dk' f(k')}{k' - (k \pm i0)}.$$

Denote by $\langle \chi, \mu \rangle_s$ a bilinear form:

$$\langle \chi, \mu \rangle_s = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} dk dl (\chi^1(l) \mu^1(k) s_2(k, l) + \chi^2(l) \mu^2(k) s_1(k, l)), \quad (3)$$

where χ^j is the element of the row χ .

Let $\varphi^{(j)}$, $j = 1, 2$ be rows which are solution of the equations conjugated to the equation for the rows $\psi^{(j)} = \{\psi_{j1}^-, \psi_{j2}^+\}$ with respect to the bilinear form (3).

Theorem 1 *Let q_1 and q_2 satisfy following conditions: $\partial^\alpha q_{1,2} \in L_1 \cap C$ at $|\alpha| \leq 3$. If $h_1(\xi, \eta)$ and $h_2(\xi, \eta)$ satisfy the conditions $\partial^\alpha h_{1,2} \in L_1 \cap C$ at $|\alpha| \leq 4$, then h_1 and h_2 may be represented in the form:*

$$\begin{aligned} h_1 &= \frac{-1}{\pi} \langle \psi^{(1)}(\xi, \eta, l), \varphi^{(1)}(\xi, \eta, k) \rangle_{\hat{h}}, \\ h_2 &= \frac{1}{\pi} \langle \psi^{(2)}(\xi, \eta, l), \varphi^{(2)}(\xi, \eta, k) \rangle_{\hat{h}}, \end{aligned} \quad (4)$$

where

$$\begin{aligned}\hat{h}_1 &= \frac{1}{4\pi}(\psi_{(1)}^+(\xi, \eta, k), \phi_{(1)}(\xi, \eta, l))_h, \\ \hat{h}_2 &= \frac{1}{4\pi}(\psi_{(2)}^-(\xi, \eta, k), \phi_{(2)}(\xi, \eta, l))_h.\end{aligned}\tag{5}$$

Of course, if $q_1 = q_2 = 0$, then the formulae (5) and (4) are ordinary Fourier transform with respect to two variables $\xi, \eta \in \mathbb{R}$.

3 The Fourier method for the linearized Davey-Stewartson I equation.

We shall consider the Davey-Stewartson I equation:

$$\begin{aligned}i\partial_t Q + (\partial_\xi^2 + \partial_\eta^2)Q + (g_1 + g_2)Q &= 0, \\ \partial_\xi g_1 = -\frac{\varepsilon}{2}\partial_\eta |Q|^2, \quad \partial_\eta g_2 = -\frac{\varepsilon}{2}\partial_\xi |Q|^2, \quad \varepsilon = \pm 1.\end{aligned}\tag{6}$$

Linearization of this equation on Q, g as a background gives:

$$\begin{aligned}i\partial_t U + (\partial_\xi^2 + \partial_\eta^2)U + (g_1 + g_2)U + (V_1 + V_2)Q &= 0, \\ \partial_\xi V_1 = -\frac{\varepsilon}{2}\partial_\eta (QU + \bar{Q}U), \quad \partial_\eta V_2 = -\frac{\varepsilon}{2}\partial_\xi (Q\bar{U} + \bar{Q}U).\end{aligned}\tag{7}$$

The equation (6) is a compatible condition for (1) at $q_1 = Q, q_2 = \varepsilon \bar{Q}$ and for a following system [7]:

$$\begin{aligned}\partial_t \psi &= i \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} (\partial_\xi - \partial_\eta)^2 \psi + i \begin{pmatrix} 0 & q_1 \\ q_2 & 0 \end{pmatrix} (\partial_\xi - \partial_\eta) \psi + \\ &+ \begin{pmatrix} ig_1 & -i\partial_\eta q_1 \\ i\partial_\xi q_2 & -ig_2 \end{pmatrix} \psi.\end{aligned}\tag{8}$$

Using the systems (1) and (8) one can prove the following statement:

Theorem 2 *Let U be the solution of (7) and Q be the solution of the Davey-Stewartson I equation with boundary conditions: $g_1|_{\xi \rightarrow \infty} = 0$ and $g_2|_{\eta \rightarrow \infty} = 0$, and U and Q satisfy the conditions of the theorem 1 for the functions h_1 and q_1 at $\forall t \in [0, T]$ respectively, then:*

$$\partial_t \hat{U}_1 = i(k^2 + l^2)\hat{U}_1, \quad \partial_t \hat{U}_2 = -i(k^2 + l^2)\hat{U}_2,\tag{9}$$

where $\hat{U}_1 = \frac{1}{4\pi}(\psi_{(1)}^-, \phi_{(1)})_U, \hat{U}_2 = \frac{1}{4\pi}(\psi_{(2)}^+, \phi_{(2)})_U$.

Inverse statement. *Let $\hat{U}_1(t, k, l)$ and $\hat{U}_2(t, k, l)$ be integrable with respect to k, l with factor $(1 + k^2)(1 + l^2)$ and satisfy equations (9), then the function*

$$U(\xi, \eta, t) = \frac{-1}{\pi} < \psi^{(1)}(\xi, \eta, l), \varphi^{(1)}(\xi, \eta, k) >_{\hat{U}}$$

satisfies linearized Davey-Stewartson I equation (7)

Solving of the system (7) is reduced to solving of the trivial equations (9) for the Fourier coefficients $\hat{U}_{1,2}$. The theorems 1 and 2 give solving (7) by Fourier method.

References

- [1] A. Davey, K. Stewartson. On three-dimensional packets of surface waves. Proc. Royal Soc. London, 1973, v.338, p.101-110.
- [2] A.S. Fokas, M.J. Ablowitz. On the inverse scattering transform of multidimensional nonlinear equations related to first-order systems in the plane. J. Math. Phys. 25(8), 1984, p.2494-2505.
- [3] A.S. Fokas, P.M.Santini. Dromions and a boundary value problem for the Davey-Stewartson I equation. Physica D, 44(1990), p.99-130.
- [4] L.P.Niznik. Inverse scattering problems for hyperbolic equations. Naukova Dumka, Kiev, 1991, 232 pp. (in russian)
- [5] D.J.Kaup. Closure of the squared Zakharov-Shabat eigenstates. J.of Math.Anal.and Appl., v.54, p.849-864, (1976).
- [6] D.J.Kaup. A perturbation expansion for the Zakharov-Shabat inverse scattering transform. SIAM J.on Appl.Math., v.31, n1, p.121-133, (1976).
- [7] M.J.Ablowitz, H.Segur. Solitons and the Inverse Scattering Transform. SIAM, Philadelphia., 1981.